

Asymptotic Behavior of the Expected Optimal Value of the Multidimensional Assignment Problem

Pavlo A. Krokmal* Don A. Grundel† Panos M. Pardalos‡

Version: August 2005

Abstract

The Multidimensional Assignment Problem (MAP) is a higher-dimensional version of the Linear Assignment Problem that arises in the areas of data association, target tracking, resource allocation, etc. This paper elucidates the question of asymptotical behavior of the expected optimal value of the large-scale MAP whose assignment costs are independent identically distributed random variables with a prescribed probability distribution. We demonstrate that for a broad class of continuous distributions the limiting value of the expected optimal cost of the MAP is determined by the location of the left endpoint of the support set of the distribution, and construct asymptotical bounds for the expected optimal cost.

Keywords: multidimensional assignment problem, random assignment problem, expected optimal value, asymptotical analysis, convergence bounds

1 Introduction

The Multidimensional Assignment Problem (MAP) is a higher dimensional version of the two-dimensional, or Linear Assignment Problem (LAP) [32]. If a classical textbook formulation of the Linear Assignment Problem is to find an optimal assignment of “ N jobs to M workers,” then, for example, the 3-dimensional Assignment Problem can be interpreted as finding an optimal assignment of “ N jobs to M workers in K time slots,” etc. In general, the objective of the MAP is to find tuples of elements from given sets, such that the total cost of the tuples is minimized. The MAP was first introduced by Pierskalla [34], and since then has found numerous applications in the areas of data association [5, 25], image recognition [39], multisensor multitarget tracking [35, 36, 24], tracking of elementary particles [38], etc. For a discussion of the MAP and its applications see, for example, [7, 9, 8] and references therein.

*Department of Mechanical and Industrial Engineering, University of Iowa, 3131 Seamans Center, Iowa City, IA 52242, E-mail: krokmal@engineering.uiowa.edu

†AFRL/MNGN, Eglin Air Force Base, E-mail: don.grundel@eglin.af.mil

‡Center for Applied Optimization, Department of Industrial and Systems Engineering, University of Florida, 303 Weil Hall, Gainesville, FL 32611, E-mail: pardalos@ufl.edu

A d -dimensional “axial” MAP with equal number n of elements in each dimension has the form

$$\begin{aligned}
\min \quad & \sum_{i_1=1}^n \cdots \sum_{i_d=1}^n c_{i_1 \dots i_d} x_{i_1 \dots i_d} & (1) \\
\text{s.t.} \quad & \sum_{i_2=1}^n \cdots \sum_{i_d=1}^n x_{i_1 \dots i_d} = 1, & i_1 = 1, \dots, n, \\
& \sum_{i_1=1}^n \cdots \sum_{i_{k-1}=1}^n \sum_{i_{k+1}=1}^n \cdots \sum_{i_d=1}^n x_{i_1 \dots i_d} = 1, & k = 2, \dots, d-1, \quad i_k = 1, \dots, n, \\
& \sum_{i_1=1}^n \cdots \sum_{i_{d-1}=1}^n x_{i_1 \dots i_d} = 1, & i_d = 1, \dots, n, \\
& x_{i_1 \dots i_d} \in \{0, 1\}^{n^d}.
\end{aligned}$$

An instance of the MAP with different numbers of elements in each dimension, $n_1 \geq n_2 \geq \dots \geq n_d$, is reducible to form (1) by introduction of dummy variables.

Problem (1) admits the following geometric interpretation: given a d -dimensional cubic matrix, find such a permutation of its rows and columns that the sum of the diagonal elements is minimized (which explains the term “axial”). This rendition leads to an alternative formulation of the MAP (1) in terms of permutations π_1, \dots, π_{d-1} of numbers 1 to n , i.e., one-to-one mappings $\pi_i : \{1, \dots, n\} \mapsto \{1, \dots, n\}$,

$$\min_{\pi_1, \dots, \pi_{d-1} \in \Pi^n} \sum_{i=1}^n c_{i, \pi_1(i), \dots, \pi_{d-1}(i)},$$

where Π^n is the set of all permutations of the set $\{1, \dots, n\}$. A feasible solution to the MAP (1) can be conveniently described by specifying its cost,

$$z = c_{i_1^{(1)} \dots i_d^{(1)}} + c_{i_1^{(2)} \dots i_d^{(2)}} + \dots + c_{i_1^{(n)} \dots i_d^{(n)}}, \quad (2)$$

where $(i_j^{(1)}, i_j^{(2)}, \dots, i_j^{(n)})$ is a permutation of the set $\{1, 2, \dots, n\}$ for every $j = 1, \dots, d$. In contrast to the LAP that represents a $d = 2$ special case of the MAP (1) and is polynomially solvable [7, 32], the MAP with $d \geq 3$ is generally NP-hard, a fact that follows from a reduction of the 3-dimensional matching problem (3DM) [14]. Although a number of exact and heuristic algorithms for the MAP [1, 6, 10, 33] have been developed in the literature, only small-to-moderate-sized instances of the MAP can be solved routinely as of today due to immense complexity of the problem.

If one denotes the optimal value of an instance of the MAP (1) with d dimensions and n elements per dimension as $z_{d,n}^*$, then it is of interest to determine the (asymptotical) behavior of $z_{d,n}^*$ when either d or n is large: $n \gg 1$ or $d \gg 1$. Evidently, such an analysis would require certain assumptions on the behavior of the assignment costs of the MAP. This is facilitated by assuming that the cost coefficients $c_{i_1 \dots i_d}$ in (1) are independent identically distributed (iid) random variables from some prescribed continuous distribution. This reduces the question at hand to asymptotical analysis of the expected optimal value $E[z_{d,n}^*]$ of random instances of the MAP, which is denoted throughout the paper as $Z_{d,n}^*$.

During the last two decades, expected optimal values of random assignment problems have been studied intensively in the context of random LAP. Perhaps, the most widely known result in this area is the

conjecture by Mézard and Parisi [22] that the expected optimal value $\mathbb{E}[L_n] := Z_{2,n}^*$ of an LAP of size n with iid uniform or exponential with mean 1 cost coefficients satisfies $\lim_{n \rightarrow \infty} \mathbb{E}[L_n] = \zeta(2) = \frac{\pi^2}{6}$, where $\zeta(\cdot)$ is Riemann's zeta function. In fact, this conjecture was preceded by an upper bound on the expected optimal value of the LAP with uniform (0,1) costs: $\limsup_{n \rightarrow \infty} L_n \leq 3$ due to Walkup [40], which was soon improved by Karp [18]: $\limsup_{n \rightarrow \infty} L_n \leq 2$. A lower bound on the limiting value of L_n was first provided by Lazarus [19, 20]: $\liminf_{n \rightarrow \infty} L_n \geq 1 + e^{-1} \approx 1.37$, and then has been improved to 1.44 by Goemans and Kodilian [15] and 1.51 by Olin [27]. Experimental evidence in support of the Mézard-Parisi conjecture was provided by Pardalos and Ramakrishnan [30]. Recently, Aldous [3] has shown that indeed $\lim_{n \rightarrow \infty} \mathbb{E}[L_n] = \frac{\pi^2}{6}$, thereby proving the conjecture (in an earlier paper [2] the author has established the existence of the limit of L_n). Another conjecture due to Parisi [31] stating that the expected optimal value of a random LAP of finite size n with exponentially distributed iid costs is equal to $\mathbb{E}[L_n] = Z_{2,n}^* = \sum_{i=1}^n i^{-2}$ has been proven independently in [21] and [26].

This paper contributes to the existing literature on random assignment problems by establishing the limiting value and asymptotic behavior of the expected optimal cost $Z_{d,n}^*$ of random Multidimensional Assignment Problem with iid cost coefficients for a broad class of continuous distributions. We demonstrate that, unlike the 2-dimensional LAP, the MAP with $d \geq 3$ has an expected optimal value that depends on the location of the left endpoint of the support set of the costs' distribution. In particular, the expected optimal value of an MAP with exponentially distributed cost coefficients approaches zero when either d or n approaches infinity. The presented analysis is constructive in the sense that it allows for deriving asymptotical lower and upper bounds for $Z_{d,n}^*$ that are converging when the support of distribution is bounded from below, as well as to estimate the rate of convergence for $Z_{d,n}^*$ when the left endpoint of the support is located at the origin.

The rest of the paper is organized as follows. The next section introduces the concept of the index tree representation of the MAP that our approach is based upon. Using the structure of the index tree graph, we construct simple expressions for the upper and lower bounds of the expected optimal cost $Z_{d,n}^*$ of the MAP. In Section 3 we establish the main result of the present endeavor by determining the limiting values of large-scale random MAPs for a wide spectrum of continuous distributions. Section 4 discusses asymptotical bounds on the expected optimal cost of random MAPs and presents the corresponding numerical results.

2 Index tree representation of the MAP and related lemmata

Our approach to determining the asymptotic behavior of the expected optimal cost $Z_{d,n}^*$ of an MAP (1) with random cost coefficients involves analysis of the so-called *index tree*, a graph structure that represents the set of feasible solutions of the MAP. First introduced by Pierskalla [34], the index tree graph $\mathcal{G} = (V, E)$ of the MAP (1) has a set of vertices V which is partitioned into n *levels*¹ and a distinct *root node*. A node at level j of the graph represents an assignment (i_1, \dots, i_d) with $i_1 = j$ and cost $c_{j i_2 \dots i_d}$, whereby each level contains $\kappa = n^{d-1}$ nodes. The set E of arcs in the index tree graph is constructed in such a way that any feasible solution of the MAP (1) can be represented as a path connecting the root node to a leaf node at level n (such a path is called a *feasible path*); evidently, the index tree contains $n!^{d-1}$ feasible paths, by the number of feasible solutions of the MAP (1).

¹In general, MAP may have n_1 elements in dimension 1, n_2 elements in dimension 2, and so on, so that in assignment (i_1, i_2, \dots, i_n) one has $1 \leq i_k \leq n_k$, $k = 1, \dots, d$. In this case, the index tree would contain n_1 levels.

The index tree representation of MAP aids in construction of lower and upper bounds for the expected optimal cost of MAP (1) with random iid costs via the following lemmata [16].

Lemma 1 *Given the index tree graph $\mathcal{G} = (V, E)$ of $d \geq 3$, $n \geq 3$ MAP whose assignment costs are iid random variables from an absolutely continuous distribution, construct set $\mathcal{A} \subset V$ by randomly selecting α different nodes from each level of the index tree. Then, \mathcal{A} is expected to contain a feasible solution of the MAP if*

$$\alpha = \left\lceil \frac{n^{d-1}}{n!^{\frac{d-1}{n}}} \right\rceil. \quad (3)$$

Lemma 2 *For a $d \geq 3$, $n \geq 3$ MAP whose cost coefficients are iid random variables from an absolutely continuous distribution F with existing first moment, define*

$$\underline{Z}_{d,n}^* := n\mathbf{E}_F [X_{(1|\kappa)}] \quad \text{and} \quad \overline{Z}_{d,n}^* := n\mathbf{E}_F [X_{(\alpha|\kappa)}], \quad (4)$$

where $X_{(i|\kappa)}$ is the i -th order statistic of $\kappa = n^{d-1}$ iid random variables with distribution F , and parameter α is determined as in (3). Then, $\underline{Z}_{d,n}^*$ and $\overline{Z}_{d,n}^*$ constitute lower and upper bounds for the expected optimal cost $Z_{d,n}^*$ of the MAP, respectively: $\underline{Z}_{d,n}^* \leq Z_{d,n}^* \leq \overline{Z}_{d,n}^*$.

Above, by an absolutely continuous distribution we mean any continuous distribution whose c.d.f. $F(\cdot)$ is representable as $F(x) = \int_{-\infty}^x f(y) dy$, where $f(\cdot)$ is the (continuous) density of the distribution.

Proofs of the lemmas are based on the probabilistic method [4] and can be found in [16]. In particular, the proof of Lemma 2 considers a set \mathcal{A}_{\min} that is constructed by selecting from each level of the index tree α nodes with the smallest costs among the κ nodes at that level. The continuity of distribution F ensures that assignment costs in the MAP (1) are all different almost surely, hence locations of the nodes that comprise the set \mathcal{A}_{\min} are random with respect to the array of nodes in each level of $\mathcal{G}(V, E)$. In the remainder of the paper, we always refer to α and κ as defined above.

As the problem size of an instance of MAP (1) is governed by two parameters, the number d of “dimensions,” and number n of elements in each dimension, asymptotic analysis of the expected optimal value $Z_{d,n}^*$ of large-scale instances of the MAP (1) involves two cases: $n \rightarrow \infty$ while d is fixed, and $d \rightarrow \infty$ for a fixed n . Throughout the paper, we use notation $\kappa \xrightarrow{n} \infty$ and $\kappa \xrightarrow{d} \infty$, respectively, to address these two situations. If a certain statement holds for both cases of $n \rightarrow \infty$ and $d \rightarrow \infty$, we indicate this by $\kappa \xrightarrow{n,d} \infty$. The rationale behind the proposed notation is that an increase in either n or d entails polynomial or exponential growth in the number κ of nodes in each level of the index tree graph: $\kappa = n^{d-1} \rightarrow \infty$.

The behavior of quantity α (3) in the cases $\kappa \xrightarrow{n} \infty$ and $\kappa \xrightarrow{d} \infty$ is more contrasting. When the number of dimensions d stays constant, and the number n of elements in each dimension increases indefinitely, α approaches a finite limiting value,

$$\alpha \rightarrow \alpha^* := \lceil e^{d-1} \rceil, \quad \kappa \xrightarrow{n} \infty, \quad (5)$$

while in the case of fixed n and unbounded d it increases exponentially:

$$\alpha \sim \kappa^{\gamma_n}, \quad \kappa \xrightarrow{d} \infty, \quad \text{where} \quad \gamma_n = 1 - \frac{\ln n!}{n \ln n}. \quad (6)$$

The range of values that parameter γ_n takes for different n is specified by the following

Lemma 3 For all $n \geq 3$, the coefficient γ_n in (6) satisfies $0 < \gamma_n < \frac{1}{2}$.

Proof: We start with establishing a simple inequality

$$n + 1 \geq \left(1 + \frac{1}{n}\right)^n, \quad n \geq 2, \quad (7)$$

that is easily validated by the induction argument. Indeed, observe that (7) holds for $n = 2$. Then, assume that (7) holds for some n , which, due to the fact that $\frac{1}{n} \geq \frac{1}{n+1}$, further implies

$$n + 1 \geq \left(1 + \frac{1}{n+1}\right)^n. \quad (8)$$

Multiplying both sides of inequality (8) by $\left(1 + \frac{1}{n+1}\right)$ we obtain

$$n + 2 = (n + 1) \left(1 + \frac{1}{n+1}\right) \geq \left(1 + \frac{1}{n+1}\right)^{n+1},$$

which verifies (7) for $n + 1$. Now, to obtain the statement of the Lemma, it is convenient to represent γ_n as

$$\gamma_n = 1 - \frac{1}{n} \sum_{r=1}^n \frac{\ln r}{\ln n}. \quad (9)$$

Evidently, since $0 < \frac{\ln r}{\ln n} < 1$ for $r = 2, \dots, n - 1$, the second term in (9) satisfies $0 < \frac{1}{n} \sum_{r=1}^n \frac{\ln r}{\ln n} < 1$. Hence, $0 < \gamma_n < 1$ for $n \geq 3$. The upper bound $\gamma_n < \frac{1}{2}$ for $n \geq 3$ is established by showing that

$$\sum_{r=1}^n \ln r > \frac{n}{2} \ln n, \quad n \geq 3. \quad (10)$$

Direct evaluation confirms that (10) holds for $n = 3$. Then, following the induction argument, we demonstrate that (9) holds for some fixed $n + 1$, provided that it holds for n . For $n + 1$ the right-hand side of (10) equals to

$$\frac{n+1}{2} \ln(n+1) = \frac{n}{2} \ln n + \frac{1}{2} \ln(n+1) + \frac{n}{2} \ln\left(1 + \frac{1}{n}\right),$$

hence, to prove that (10) holds for $n + 1$ it suffices to show that

$$\ln(n+1) \geq \frac{1}{2} \ln(n+1) + \frac{n}{2} \ln\left(1 + \frac{1}{n}\right).$$

The last inequality is equivalent to (7), which proves the Lemma. ■

The reader has already noticed that the presented lemmata addresses only MAPs with $d \geq 3$, $n \geq 3$. Before proceeding further, we comment shortly on two special cases of the MAP (1) when $d = 2$ or $n = 2$. The case $d = 2$ represents, as noted earlier, the Linear Assignment Problem, whose asymptotic behavior is distinctly different from that of MAPs with $d \geq 3$ (see Sections 3 and 4). It can be shown that in the case of $d = 2$ Lemmas 1 and 2 produce only trivial bounds that are rather inefficient in determining the asymptotic behavior of the expected optimal value of the LAP within the presented approach.

When $n = 2$, the costs of feasible solutions to the MAP (1) have the form

$$z = c_{i_1^{(1)} \dots i_d^{(1)}} + c_{i_1^{(2)} \dots i_d^{(2)}}, \quad \text{where } i_j^{(1)}, i_j^{(2)} \in \{1, 2\}, \quad i_j^{(1)} \neq i_j^{(2)},$$

and consequently are iid random variables with distribution F_2 , which is the convolution of F with itself: $F_2 = F * F$ [17]. Then, obviously, the expected optimal value of a random MAP with $n = 2$ can be expressed as expectation of the minimum order statistic of 2^{d-1} iid variables with distribution F_2 :

$$Z_{d,2}^* = \mathbb{E}_{F * F} [X_{(1|2^{d-1})}], \quad (11)$$

which simplifies the analysis of the expected optimal cost as compared to the general case of $n \geq 3$ that necessitates the use of the lower and upper bounds (4).

3 Main result

In this section we derive the limiting values of the expected optimal cost $Z_{d,n}^*$ of a random MAP (1) with iid cost coefficients drawn from a broad class of continuous distributions, when one of the dimension parameters n or d of the problem increases indefinitely. First we lay out preliminary considerations that make a foundation for more intricate analysis presented below.

According to Lemma 2, the upper bound on the expected optimal cost $Z_{d,n}^*$ of a random MAP with iid cost coefficients from an absolutely continuous distribution F is given by $\bar{Z}_{d,n}^* = n \mathbb{E}_F [X_{(\alpha|\kappa)}]$, where $X_{(\alpha|\kappa)}$ is the α -th order statistic among κ independent F -distributed random variables. It is well-known (see, e.g., [11]) that order statistics have distributions of beta type; for instance, the distribution of $X_{(\alpha|\kappa)}$ is given by

$$\mathbb{P}[X_{(\alpha|\kappa)} \leq x] = \frac{\kappa!}{(\alpha-1)!(\kappa-\alpha)!} \int_0^{F(x)} t^{\alpha-1} (1-t)^{\kappa-\alpha} dt.$$

Then, assuming that the interior of the support set of distribution F has the form (a, b) , where

$$a = \inf \{x \mid F(x) > 0\}, \quad b = \sup \{x \mid F(x) < 1\},$$

the upper bound $\bar{Z}_{d,n}^*$ can be represented as

$$\bar{Z}_{d,n}^* = \frac{n \Gamma(\kappa + 1)}{\Gamma(\alpha) \Gamma(\kappa - \alpha + 1)} I_{\alpha,\kappa}, \quad (12)$$

with $I_{\alpha,\kappa}$ being the integral of the form

$$I_{\alpha,\kappa} = \int_a^b x [F(x)]^{\alpha-1} [1 - F(x)]^{\kappa-\alpha} dF(x) = \int_0^1 F^{-1}(u) u^{\alpha-1} (1-u)^{\kappa-\alpha} du. \quad (13)$$

In the last equality, $F^{-1}(\cdot)$ denotes the inverse of the c.d.f. $F(x)$ of distribution of assignment costs in the MAP (1). While it is practically impossible to evaluate integral (13) exactly in the general case, the asymptotic behavior of (13) for large n and d can be determined for a wide range of distributions F . In particular, it will be seen that the asymptotic value of $I_{\alpha,\kappa}$, and, consequently, that of $Z_{d,n}^*$ for $n, d \gg 1$ depends on the location of the left endpoint a of the support of F .

The exposition of the results in the sequel assumes familiarity with the foundations of asymptotical analysis and theory of special functions. For example, the asymptotic properties of integrals of type (13) for large values of n and d can be quantified in terms of the Beta and Gamma functions,

$$\mathbb{B}(x, y) = \int_0^1 t^{x-1} (1-t)^{y-1} dt = \frac{\Gamma(x) \Gamma(y)}{\Gamma(x+y)}, \quad \Gamma(z) = \int_0^\infty e^{-t} t^{z-1} dt. \quad (14)$$

Although we make extensive use of these and other related special functions, we do not discuss herein their numerous properties that are used to derive the majority of the results of this paper. Instead, we refer the reader to an excellent treatise on special functions edited by Erdélyi [13]; also, a comprehensive collection of facts and formulae on special functions is available at [41]. Text by Olver [28] is another excellent reference to both asymptotical analysis and the theory of special functions.

3.1 Distributions with a finite left endpoint of the support

The asymptotic analysis of the expected optimal value of the random MAPs with continuous distributions is facilitated by means of the following lemma, which provides asymptotic series representations on a scale of Beta functions for integrals of type (13).

Lemma 4 *Let function $h(u)$ have the following asymptotic expansion at $0+$,*

$$h(u) \sim \sum_{s=0}^{\infty} a_s u^{(s+\lambda-\mu)/\mu}, \quad u \rightarrow 0+, \quad (15)$$

where $\lambda, \mu > 0$. Then for any positive integer m one has

$$\int_0^1 h(u) u^{\alpha-1} (1-u)^{\kappa-\alpha} du = \sum_{s=0}^{m-1} a_s \phi_s(\kappa) + \mathcal{O}(\phi_m(\kappa)), \quad \kappa \xrightarrow{n,d} \infty, \quad (16)$$

where $\phi_s(\kappa) = \mathbf{B}\left(\frac{s+\lambda}{\mu} + \alpha - 1, \kappa - \alpha + 1\right)$, $s = 0, 1, \dots$, provided that the integral is absolutely convergent for $\kappa = \alpha = 1$.

Proof: First we demonstrate that the sequence of functions $\{\phi_s(\kappa)\}_{s=0}^{\infty}$ represents a *scale* [28] in both cases of large n and d , i.e., $\phi_{s+1}(\kappa) = o(\phi_s(\kappa))$, $\kappa \xrightarrow{n,d} \infty$. The ratio of $\phi_{s+1}(\kappa)$ and $\phi_s(\kappa)$ has the form

$$\frac{\phi_{s+1}(\kappa)}{\phi_s(\kappa)} = \frac{\Gamma\left(\frac{s+1+\lambda}{\mu} + \alpha - 1\right)}{\Gamma\left(\frac{s+\lambda}{\mu} + \alpha - 1\right)} \cdot \frac{\Gamma\left(\kappa + \frac{s+\lambda}{\mu}\right)}{\Gamma\left(\kappa + \frac{s+1+\lambda}{\mu}\right)}.$$

Using a well-known expansion for the quotient of the Gamma functions [13]

$$\frac{\Gamma(z+a)}{\Gamma(z+b)} = z^{a-b} \left(1 + \mathcal{O}(z^{-1})\right), \quad z \rightarrow \infty, \quad a, b \in \mathbb{R}, \quad (17)$$

we obtain that for a fixed d and $n \rightarrow \infty$,

$$\frac{\phi_{s+1}(\kappa)}{\phi_s(\kappa)} \sim \frac{\Gamma\left(\frac{s+1+\lambda}{\mu} + \alpha^* - 1\right)}{\Gamma\left(\frac{s+\lambda}{\mu} + \alpha^* - 1\right)} \kappa^{-1/\mu} \rightarrow 0, \quad \kappa \xrightarrow{n} \infty, \quad \mu > 0.$$

Similarly, for $d \rightarrow \infty$ and n fixed, by virtue of Lemma 3 one has

$$\frac{\phi_{s+1}(\kappa)}{\phi_s(\kappa)} \sim \frac{\kappa^{\gamma n/\mu}}{\kappa^{1/\mu}} = \kappa^{\frac{\gamma n-1}{\mu}} \rightarrow 0, \quad \kappa \xrightarrow{d} \infty, \quad \mu > 0.$$

Now we show that expansion (16) holds. For any non-negative integer m , let

$$h(u) = \sum_{s=0}^{m-1} a_s u^{(s+\lambda-\mu)/\mu} + \varphi_m(u), \quad 0 < u < 1. \quad (18)$$

Substituting the above equality in the left-hand side of (16), one obtains

$$\begin{aligned} & \int_0^1 h(u) u^{\alpha-1} (1-u)^{\kappa-\alpha} du \\ &= \sum_{s=0}^{m-1} a_s \mathbf{B}\left(\frac{s+\lambda}{\mu} + \alpha - 1, \kappa - \alpha + 1\right) + \int_0^1 \varphi_m(u) u^{\alpha-1} (1-u)^{\kappa-\alpha} du. \end{aligned} \quad (19)$$

To estimate the remainder term in (19), observe that $\varphi_m(u) = \mathcal{O}(u^{(m+\lambda-\mu)/\mu})$ for $u \rightarrow 0+$, in accordance with the definition of asymptotic expansion (15). Consequently, there exist $u_m \in (0, 1)$ such that $C_m = \sup_{u \in (0, u_m]} |\varphi_m(u) u^{-(m+\lambda-\mu)/\mu}|$ is finite, whence

$$\left| \int_0^{u_m} \varphi_m(u) u^{\alpha-1} (1-u)^{\kappa-\alpha} du \right| \leq C_m \mathbf{B}\left(\frac{m+\lambda}{\mu} + \alpha - 1, \kappa - \alpha + 1\right). \quad (20)$$

Since integral (16) is absolutely convergent for $\kappa = \alpha = 1$, the integral $\int_{u_m}^1 \varphi_m(u) du$ is absolutely convergent too, which implies absolute convergence of $\int_{u_m}^1 \varphi_m(u) u^{-M} du$ for any $M > 0$:

$$\int_{u_m}^1 |\varphi_m(u)| du \leq \int_{u_m}^1 |\varphi_m(u) u^{-M}| du \leq u_m^{-M} \int_{u_m}^1 |\varphi_m(u)| du.$$

For $M = \frac{m+\lambda}{\mu} + 1$ denote $\Xi_m = \sup_{\tau \in [u_m, 1]} |\zeta_m(\tau)|$, where $\zeta_m(\tau) = \int_{u_m}^{\tau} \varphi_m(u) u^{-M} du$. Integrating the remainder term in (19) by parts, we have

$$\begin{aligned} \int_{u_m}^1 \varphi_m(u) u^{\alpha-1} (1-u)^{\kappa-\alpha} du &= -(\alpha + M - 1) \int_{u_m}^1 \zeta_m(u) u^{\alpha+M-2} (1-u)^{\kappa-\alpha} du \\ &\quad + (\kappa - \alpha) \int_{u_m}^1 \zeta_m(u) u^{\alpha+M-1} (1-u)^{\kappa-\alpha-1} du, \end{aligned} \quad (21)$$

which implies that the first integral in (21) is bounded as

$$\left| \int_{u_m}^1 \varphi_m(u) u^{\alpha-1} (1-u)^{\kappa-\alpha} du \right| < 2 \Xi_m \frac{\Gamma(\alpha + M) \Gamma(\kappa - \alpha + 1)}{\Gamma(\kappa + M)}. \quad (22)$$

Considering the behavior of the right-hand side of (22) with respect to $\phi_m(\kappa)$ for large κ ,

$$\frac{\Gamma(\alpha + M) \Gamma(\kappa - \alpha + 1)}{\Gamma(\kappa + M)} \cdot \frac{\Gamma\left(\kappa + \frac{m+\lambda}{\mu}\right)}{\Gamma\left(\alpha + \frac{m+\lambda}{\mu} - 1\right) \Gamma(\kappa - \alpha + 1)},$$

we see that their ratio is of order of κ^{-1} when $\kappa \xrightarrow{n} \infty$, and is $\mathcal{O}(\kappa^{2\gamma_n-1})$ if $\kappa \xrightarrow{d} \infty$. Hence, by Lemma 3 we have that

$$\left| \int_{u_m}^1 \varphi_m(u) u^{\alpha-1} (1-u)^{\kappa-\alpha} du \right| = o\left(\mathbf{B}\left(\frac{m+\lambda}{\mu} + \alpha - 1, \kappa - \alpha + 1\right)\right), \quad \kappa \xrightarrow{n,d} \infty. \quad (23)$$

Combining (19), (20), and (23) we arrive at the sought equality (16). ■

With the help of Lemma 4 it is now straightforward to derive the limiting value of the expected optimal cost of MAP with random iid cost coefficients that have a continuous p.d.f. with a finite left endpoint of the support.

Theorem 1 Consider a $d \geq 3$, $n \geq 3$ MAP (1) with cost coefficients that are iid random variables from an absolutely continuous distribution with existing first moment. Let (a, b) , where $a \in \mathbb{R}$, $b \in (a, +\infty]$, be the interior of the support set of this distribution. Further, assume that the inverse $F^{-1}(x)$ of the c.d.f. $F(x)$ of the distribution is such that

$$F^{-1}(x) = a + \mathcal{O}(x^\beta), \quad x \rightarrow 0+, \quad \beta > 0. \quad (24)$$

Then, for a fixed n and $d \rightarrow \infty$, or a fixed d and $n \rightarrow \infty$, the expected optimal value $Z_{d,n}^*$ satisfies

$$\lim_{d \rightarrow \infty} Z_{d,n}^* = na, \quad \lim_{n \rightarrow \infty} Z_{d,n}^* = \begin{cases} +\infty, & a > 0, \\ 0, & a = 0, \\ -\infty, & a < 0. \end{cases} \quad (25)$$

Proof: According to the aforesaid, an upper bound on the expected optimal value of a random MAP whose assignment costs are iid random variables with distribution F is obtained via (12)–(13). Existence of the first moment of F ensures the existence of its order statistics, as well as absolute integrability of $F^{-1}(\cdot)$ on the interval $(0, 1)$. Given the asymptotic behavior (24) of function $F^{-1}(u)$ at $u = 0+$, application of Lemma 4 to integral $I_{a,\kappa}$ (13) produces

$$I_{a,\kappa} = a \frac{\Gamma(\alpha) \Gamma(\kappa - \alpha + 1)}{\Gamma(\kappa + 1)} + \mathcal{O}\left(\frac{\Gamma(\alpha + \beta) \Gamma(\kappa - \alpha + 1)}{\Gamma(\kappa + \beta + 1)}\right), \quad \kappa \xrightarrow{n,d} \infty,$$

which, upon substitution into (12), leads to

$$\bar{Z}_{d,n}^* = na + \mathcal{O}\left(n \frac{\Gamma(\alpha + \beta) \Gamma(\kappa + 1)}{\Gamma(\alpha) \Gamma(\kappa + \beta + 1)}\right), \quad \kappa \xrightarrow{n,d} \infty. \quad (26)$$

By asymptotic expansion (17) the second term in (26) vanishes as $\kappa \xrightarrow{n,d} \infty$, yielding that the upper bound $\bar{Z}_{d,n}^*$ satisfies the relations (25). On the other hand, since a is the left endpoint of the support set of distribution F , the cost z of a feasible solution (2) of the MAP is bounded from below by $\inf z = na$. It is easy to see that $\inf z$ does also satisfy equalities (25), which, by virtue of inequality

$$\inf z \leq Z_{d,n}^* \leq \bar{Z}_{d,n}^*,$$

proves the statement (25) of the Theorem. ■

The conditions of Theorem 1 are quite general: it is only required that function $F^{-1}(\cdot)$ is absolutely integrable on $(0, 1)$ and there exists $\beta > 0$ such that

$$\lim_{x \rightarrow 0+} |x^{-\beta} (F^{-1}(x) - a)| < \infty.$$

The case when distribution F is such that the rate of convergence of $F^{-1}(x) - a$ as $x \rightarrow 0+$ is slower than that of x^β , for some $\beta > 0$, can be considered similarly, based on the particular form of $F^{-1}(\cdot)$.

The proved theorem evidences that the asymptotic behavior of the expected optimal value of a random MAP whose distribution has a bounded from below support is quite dissimilar in cases of $d \gg 1$ and $n \gg 1$. Indeed, for a fixed d and $n \rightarrow \infty$, the expected optimal value $Z_{d,n}^*$ of a random MAP with corresponding distribution F is finite only when the left endpoint of the support of F is located at the origin: $a = 0$. On the contrary, $\lim_{d \rightarrow \infty} Z_{d,n}^*$ for a fixed n is finite as long as a is finite.

Trivially, finiteness of the left endpoint a of the support of the distribution implies finiteness of the function $F^{-1}(\cdot)$ at the origin: $F^{-1}(0) = a \in \mathbb{R}$. If the support of distribution F has infinite left endpoint, $a = -\infty$, the inverse $F^{-1}(x)$ of the distribution's c.d.f. necessarily has a singularity at $x = 0+$:

$$\lim_{x \rightarrow 0+} F^{-1}(x) = -\infty.$$

The next subsection contains an analysis of the expected optimal value of MAP with iid cost coefficients drawn from a continuous distribution whose support has infinite left endpoint.

3.2 Distributions with infinite left endpoint of the support

The case of a continuous distribution with infinite left endpoint of the support set is more cumbersome as different singularities of $F^{-1}(x)$ at $x = 0+$ require differing approaches to their treatment. To this end, we consider two most common singularities of $F^{-1}(0+)$ encountered in applications: a power singularity and a logarithmic singularity.

We say that $F^{-1}(x)$ has a power singularity at $x = 0+$ if $F^{-1}(x) \sim -\nu x^{-\beta}$, $x \rightarrow 0+$, where ν and β are positive constants. It is easy to see that for function $F^{-1}(x)$ to be integrable over the interval $(0, 1)$, the range of values for parameter β must be limited to $0 < \beta < 1$. The next theorem addresses this case.

Theorem 2 Consider a $d \geq 3$, $n \geq 3$ MAP (1) with cost coefficients that are iid random variables from an absolutely continuous distribution whose expectation exists. Let $(-\infty, b)$, where $b \in (-\infty, +\infty]$, be the support set of this distribution. Further, assume that the inverse $F^{-1}(x)$ of the c.d.f. $F(x)$ of the distribution has an asymptotic expansion at $x \rightarrow 0+$

$$F^{-1}(x) \sim -\nu x^{-\beta}, \quad x \rightarrow 0+, \quad 0 < \beta < 1, \quad \nu > 0. \quad (27)$$

Then, for a fixed n and $d \rightarrow \infty$, or a fixed d and $n \rightarrow \infty$, the expected optimal value $Z_{d,n}^*$ of the MAP is unbounded from below:

$$\lim_{d \rightarrow \infty} Z_{d,n}^* = \lim_{n \rightarrow \infty} Z_{d,n}^* = -\infty. \quad (28)$$

Proof: Essentially the same as that of Theorem 1. ■

Now we examine the case of the inverse $F^{-1}(x)$ of distribution $F(\cdot)$ having a (weaker) logarithmic singularity at zero, $F^{-1}(x) \sim -\nu (\ln \frac{1}{x})^\beta$, $x \rightarrow 0+$, where ν and β are again positive numbers. Similarly, in this case the expected optimal value of MAP is also unbounded from below. A key element of the subsequent analysis is the asymptotical valuation of the integral

$$\int_0^1 u^{\alpha+\delta-1} (1-u)^{\kappa-\alpha} \ln \frac{1}{u} du, \quad (29)$$

where $\delta > 0$, for large values of n and d . By differentiating the identity for the Beta function (see (14))

$$\int_0^1 u^{\alpha+\delta-1}(1-u)^{\kappa-\alpha} du = \frac{\Gamma(\alpha+\delta)\Gamma(\kappa-\alpha+1)}{\Gamma(\kappa+\delta+1)}$$

with respect to α , one has

$$\begin{aligned} \int_0^1 u^{\alpha+\delta-1}(1-u)^{\kappa-\alpha} \ln \frac{1}{u} du &= -\frac{1}{\Gamma(\kappa+\delta+1)} \frac{\partial}{\partial \alpha} [\Gamma(\alpha+\delta)\Gamma(\kappa-\alpha+1)] \\ &\quad - \int_0^1 u^{\alpha+\delta}(1-u)^{\kappa-\alpha} \ln(1-u) du. \end{aligned} \quad (30)$$

Given that $\ln(1-u) = -u + \mathcal{O}(u^2)$, $u \rightarrow 0+$, the asymptotic value of the last integral in (30) is obtained by application of Lemma 4 as

$$\int_0^1 u^{\alpha+\delta}(1-u)^{\kappa-\alpha} \ln(1-u) du = \mathcal{O}\left(\frac{\Gamma(\alpha+\delta+1)\Gamma(\kappa-\alpha+1)}{\Gamma(\kappa+\delta+2)}\right).$$

Next, the first term in the right-hand side of (30) can be rewritten using the logarithmic derivative of the Gamma function $\psi(z) = \Gamma'(z)/\Gamma(z)$ in the form

$$\frac{\Gamma(\alpha+\delta)\Gamma(\kappa-\alpha+1)}{\Gamma(\kappa+\delta+1)} [\psi(\kappa-\alpha+1) - \psi(\alpha+\delta)].$$

Employing the asymptotic expansion of $\psi(z)$ for large values of argument [13],

$$\psi(z) = \ln z - \frac{1}{2z} + \mathcal{O}(z^{-2}), \quad z \rightarrow \infty, \quad (31)$$

we arrive at the sought asymptotical expressions for integral (29). In the case $n \rightarrow \infty$, we have

$$\int_0^1 u^{\alpha+\delta-1}(1-u)^{\kappa-\alpha} \ln \frac{1}{u} du = \frac{\Gamma(\alpha^*+\delta)}{\kappa^{\alpha^*+\delta}} \ln \kappa \left(1 + \mathcal{O}\left(\frac{1}{\ln \kappa}\right)\right), \quad \kappa \xrightarrow{n} \infty. \quad (32)$$

In the case of $d \rightarrow \infty$, integral (29) equals

$$\int_0^1 u^{\alpha+\delta-1}(1-u)^{\kappa-\alpha} \ln \frac{1}{u} du = \frac{\Gamma(\kappa^{\gamma_n} + \delta)\Gamma(\kappa - \kappa^{\gamma_n} + 1)}{\Gamma(\kappa + \delta + 1)} \left\{ (1 - \gamma_n) \ln \kappa + \mathcal{O}(\kappa^{-\gamma_n}) \right\}, \quad \kappa \xrightarrow{d} \infty. \quad (33)$$

By use of the expansion

$$\frac{\Gamma(\kappa^{\gamma_n} + \delta)}{\Gamma(\kappa + \delta + 1)} = \frac{e^{\kappa - \kappa^{\gamma_n}} \kappa^{(1-3\gamma_n)/2}}{\kappa^{\kappa - \gamma_n \kappa^{\gamma_n}}} \kappa^{-(1-\gamma_n)(\delta+1)} \left(1 + \mathcal{O}(\kappa^{-\gamma_n})\right), \quad \kappa \xrightarrow{d} \infty, \quad (34)$$

expression (33) can be transformed to

$$\int_0^1 u^{\alpha+\delta-1}(1-u)^{\kappa-\alpha} \ln \frac{1}{u} du = R_{\gamma_n}^{\kappa} \kappa^{-(1-\gamma_n)(\delta+1)} (1 - \gamma_n) \ln \kappa \left(1 + \mathcal{O}(\kappa^{-\gamma_n})\right), \quad \kappa \xrightarrow{d} \infty, \quad (35)$$

where $R_{\gamma_n}^{\kappa}$ stands for

$$R_{\gamma_n}^{\kappa} = \Gamma(\kappa - \kappa^{\gamma_n} + 1) e^{\kappa - \kappa^{\gamma_n}} \kappa^{-\kappa + \gamma_n \kappa^{\gamma_n} + (1-3\gamma_n)/2}. \quad (36)$$

Now we are ready to prove the main result of this subsection.

Theorem 3 Consider a $d \geq 3$, $n \geq 3$ MAP (1) with cost coefficients that are iid random variables from an absolutely continuous distribution F whose expectation exists. Let $(-\infty, b)$, where $b \in (-\infty, +\infty]$, be the interior of the support set of F , and assume that the inverse $F^{-1}(x)$ of the c.d.f. $F(x)$ of the distribution has a logarithmic singularity at $x \rightarrow 0+$,

$$F^{-1}(x) \sim -\nu \left(\ln \frac{1}{x}\right)^\beta, \quad x \rightarrow 0+, \quad \text{where } \beta > 0, \quad \nu > 0. \quad (37)$$

Then, for a fixed n and $d \rightarrow \infty$, or a fixed d and $n \rightarrow \infty$, the expected optimal value $Z_{d,n}^*$ of the MAP is unbounded from below:

$$\lim_{d \rightarrow \infty} Z_{d,n}^* = \lim_{n \rightarrow \infty} Z_{d,n}^* = -\infty. \quad (38)$$

Proof: As before, the proof is based on examining the asymptotical behavior of the upper bound $\bar{Z}_{d,n}^*$, which in the case of a logarithmic singularity (37) of $F^{-1}(x)$ at $x = 0+$ comes down to asymptotic evaluation of the integral

$$\int_0^1 u^{\alpha-1} (1-u)^{\kappa-\alpha} \left(\ln \frac{1}{u}\right)^\beta du \quad (39)$$

for large values of n and d . The asymptotical properties of this integral as n or d approach infinity are determined by the behavior of the integrand in the vicinity of 0; in particular, for any $\beta > 0$ one has

$$\int_0^1 u^{\alpha-1} (1-u)^{\kappa-\alpha} \left(\ln \frac{1}{u}\right)^\beta du = \int_0^{1/e} u^{\alpha-1} (1-u)^{\kappa-\alpha} \left(\ln \frac{1}{u}\right)^\beta du + \mathcal{O}\left(\frac{\Gamma(\alpha) \Gamma(\kappa - \alpha + 1)}{\Gamma(\kappa + 1)}\right), \quad (40)$$

for $\kappa \xrightarrow{n,d} \infty$. Indeed, observe that $\frac{1}{e} \leq u \leq 1$ implies that $0 \leq \left(\ln \frac{1}{u}\right)^\beta \leq 1$ for $\beta > 0$, therefore the following integral is bounded as

$$\int_{1/e}^1 u^{\alpha-1} (1-u)^{\kappa-\alpha} \left(\ln \frac{1}{u}\right)^\beta du \leq \int_{1/e}^1 u^{\alpha-1} (1-u)^{\kappa-\alpha} du \leq \int_0^1 u^{\alpha-1} (1-u)^{\kappa-\alpha} du = B(\alpha, \kappa - \alpha + 1),$$

which entails the following asymptotic expression that verifies equality (40):

$$\int_{1/e}^1 u^{\alpha-1} (1-u)^{\kappa-\alpha} \left(\ln \frac{1}{u}\right)^\beta du = \mathcal{O}\left(\frac{\Gamma(\alpha) \Gamma(\kappa - \alpha + 1)}{\Gamma(\kappa + 1)}\right), \quad \kappa \xrightarrow{n,d} \infty. \quad (41)$$

The choice of constant $1/e$ that splits the integration interval in (40) is merely due to our convenience; a similar decomposition can be obtained for any small enough positive number between 0 and 1. Following [12], to estimate the asymptotical behavior of integral

$$\int_0^{1/e} u^{\alpha-1} (1-u)^{\kappa-\alpha} \left(\ln \frac{1}{u}\right)^\beta du := J(\alpha, \kappa - \alpha + 1, \beta), \quad (42)$$

that enters the right-hand side of decomposition (40), we employ the functional equation

$$\frac{1}{\Gamma(\varrho)} \int_0^\infty z^{\varrho-1} J(\alpha + z, \kappa, \beta) dz = J(\alpha, \kappa, \beta - \varrho), \quad \alpha, \beta, \kappa, \varrho > 0, \quad (43)$$

which is easily derivable by plugging (42) into the left-hand side of (43) and interchanging the order of integration. First we consider the case of $0 < \beta < 1$. For $\varrho \in (0, 1)$ let $\beta = 1 - \varrho$, then integral $J(\alpha, \kappa - \alpha + 1, \beta)$ (42) can be written as a sum of two other integrals, J_1 and J_2 ,

$$J(\alpha, \kappa - \alpha + 1, 1 - \varrho) = \frac{1}{\Gamma(\varrho)} \left(\int_0^\zeta + \int_\zeta^\infty \right) z^{\varrho-1} J(\alpha + z, \kappa - \alpha + 1, 1) dz = J_1 + J_2, \quad (44)$$

where $\zeta > 0$ is a positive number. With interchange of the order of integration, J_2 reduces to

$$J_2 = \frac{1}{\Gamma(\varrho)} \int_0^{1/e} u^{\alpha+\zeta} (1-u)^{\kappa-\alpha} \ln \frac{1}{u} du \int_\zeta^\infty z^{\varrho-1} u^{z-\zeta} dz. \quad (45)$$

Since $0 \leq u \leq \frac{1}{e}$, the inner integral in (45) is a bounded function of u ,

$$\int_\zeta^\infty z^{\varrho-1} u^{z-\zeta} dz = \int_\zeta^\infty z^{\varrho-1} e^{-(z-\zeta) \ln u} dz \leq \int_\zeta^\infty z^{\varrho-1} e^{-(z-\zeta)} dz,$$

hence, J_2 can be estimated as

$$J_2 = \mathcal{O} \left(\int_0^{1/e} u^{\alpha+\zeta-1} (1-u)^{\kappa-\alpha} \ln \frac{1}{u} du \right), \quad \kappa \xrightarrow{n,d} \infty. \quad (46)$$

Next, consider integral J_1 in (44):

$$J_1 = \frac{1}{\Gamma(\varrho)} \int_0^\zeta z^{\varrho-1} dz \int_0^{1/e} u^{\alpha+z-1} (1-u)^{\kappa-\alpha} \ln \frac{1}{u} du. \quad (47)$$

By means of decomposition (40), the inner integral in (47) can be represented as

$$\begin{aligned} & \int_0^{1/e} u^{\alpha+z-1} (1-u)^{\kappa-\alpha} \ln \frac{1}{u} du \\ &= \int_0^1 u^{\alpha+z-1} (1-u)^{\kappa-\alpha} \ln \frac{1}{u} du + \mathcal{O} \left(\frac{\Gamma(\alpha+z) \Gamma(\kappa-\alpha+1)}{\Gamma(\kappa+z+1)} \right), \quad \kappa \xrightarrow{n,d} \infty. \end{aligned} \quad (48)$$

Further treatment of integrals J_1 and J_2 depends on the rate with which κ approaches infinity, i.e., which of the parameters n and d is fixed and which is infinitely increasing. We structure the remainder of the proof of the theorem in the following steps.

Step 1: $\kappa \xrightarrow{n} \infty$. If n goes to infinity and d is fixed, we have by (5) that $\alpha \rightarrow \alpha^* \in \mathbb{R}$. In accordance to the derived asymptotic expansion (32), the integral in the right-hand side of (48) equals

$$\int_0^1 u^{\alpha+z-1} (1-u)^{\kappa-\alpha} \ln \frac{1}{u} du = \frac{\Gamma(\alpha^*+z)}{\kappa^{\alpha^*+z}} \ln \kappa \left(1 + \mathcal{O} \left(\frac{1}{\ln \kappa} \right) \right), \quad \kappa \xrightarrow{n} \infty.$$

Similarly, the last term in (48) admits the representation

$$\frac{\Gamma(\alpha+z) \Gamma(\kappa-\alpha+1)}{\Gamma(\kappa+z+1)} = \mathcal{O} \left(\kappa^{-\alpha^*-z} \right), \quad \kappa \xrightarrow{n} \infty.$$

On inserting the last two equalities in (48) we obtain

$$J_1 = \frac{1}{\Gamma(\varrho) \kappa^{\alpha^*}} \left\{ \left(\ln \kappa + \mathcal{O}(1) \right) \int_0^\zeta z^{\varrho-1} \Gamma(\alpha^*+z) \kappa^{-z} dz + \int_0^\zeta z^{\varrho-1} \mathcal{O}(\kappa^{-z}) dz \right\}. \quad (49)$$

Now, consider the first integral in the right-hand side of (49). Since $\Gamma(\alpha^* + z) = \Gamma(\alpha^*) + \mathcal{O}(z)$, $z \rightarrow 0+$, this integral can be restated as

$$\int_0^\zeta z^{\varrho-1} \Gamma(\alpha^* + z) \kappa^{-z} dz = \Gamma(\alpha^*) \int_0^\zeta z^{\varrho-1} \kappa^{-z} dz + \mathcal{O}\left(\int_0^\zeta z^\varrho \kappa^{-z} dz\right). \quad (50)$$

The first integral in the right-hand side of (50) can be expressed in terms of the incomplete Gamma function $\Gamma(a, \zeta) = \int_\zeta^\infty e^{-t} t^{a-1} dt$ [13]:

$$\int_0^\zeta z^{\varrho-1} \kappa^{-z} dz = \frac{\Gamma(\varrho)}{(\ln \kappa)^\varrho} - \frac{\Gamma(\varrho, \zeta \ln \kappa)}{(\ln \kappa)^\varrho}. \quad (51)$$

Using the following asymptotic relation for the incomplete Gamma function [13]

$$\frac{\Gamma(a, \zeta x)}{x^a} = \mathcal{O}\left(\frac{e^{-\zeta x}}{x}\right), \quad x \rightarrow \infty, \quad a > 0, \quad (52)$$

the integrals in (50) can be evaluated as

$$\int_0^\zeta \kappa^{-z} z^{\varrho-1} dz = \frac{\Gamma(\varrho)}{(\ln \kappa)^\varrho} + \mathcal{O}\left(\frac{1}{\kappa^\zeta \ln \kappa}\right), \quad \mathcal{O}\left(\int_0^\zeta z^\varrho \kappa^{-z} dz\right) = \mathcal{O}\left((\ln \kappa)^{-\varrho-1}\right), \quad \kappa \xrightarrow{n} \infty,$$

which yields the asymptotic value of (50) in the form

$$\int_0^\zeta z^{\varrho-1} \Gamma(\alpha^* + z) \kappa^{-z} dz = \frac{\Gamma(\alpha^*) \Gamma(\varrho)}{(\ln \kappa)^\varrho} \left(1 + \mathcal{O}\left(\frac{1}{\ln \kappa}\right)\right), \quad \kappa \xrightarrow{n} \infty.$$

On substitution of the last expression for integral (50) into representation (49) for J_1 we have

$$J_1 = \frac{\Gamma(\alpha^*)}{\kappa^{\alpha^*}} (\ln \kappa)^\beta \left(1 + \mathcal{O}\left(\frac{1}{\ln \kappa}\right)\right), \quad \kappa \xrightarrow{n} \infty,$$

and, invoking expansions (46) and (32) with ζ large enough, we obtain

$$\int_0^1 u^{\alpha-1} (1-u)^{\kappa-\alpha} \left(\ln \frac{1}{u}\right)^\beta du = \frac{\Gamma(\alpha^*)}{\kappa^{\alpha^*}} (\ln \kappa)^\beta \left(1 + \mathcal{O}\left(\frac{1}{(\ln \kappa)^\beta}\right)\right), \quad \kappa \xrightarrow{n} \infty. \quad (53)$$

To derive the counterpart of equality (53) for the case $\kappa \xrightarrow{d} \infty$, one has to take a slightly different route.

Step 2: $\kappa \xrightarrow{d} \infty$. In the case of d increasing and n fixed, we have that both α and κ are increasing, such that $\alpha \sim \kappa^{\gamma_n}$, $\kappa \rightarrow \infty$, where $0 < \gamma_n < \frac{1}{2}$, according to (6) and Lemma 3. By virtue of representation (35), the integral in the right-hand side of (48) has the form

$$\int_0^1 u^{\kappa^{\gamma_n} + z - 1} (1-u)^{\kappa - \kappa^{\gamma_n}} \ln \frac{1}{u} du = R_{\gamma_n}^\kappa \kappa^{-(1-\gamma_n)(z+1)} (1-\gamma_n) \ln \kappa \left(1 + \mathcal{O}(\kappa^{-\gamma_n})\right), \quad \kappa \xrightarrow{d} \infty.$$

Similarly, (34) yields

$$\mathcal{O}\left(\frac{\Gamma(\kappa^{\gamma_n} + z) \Gamma(\kappa - \kappa^{\gamma_n} + 1)}{\Gamma(\kappa + z + 1)}\right) = R_{\gamma_n}^\kappa \kappa^{-(1-\gamma_n)} \mathcal{O}\left(\kappa^{-(1-\gamma_n)z}\right), \quad \kappa \xrightarrow{d} \infty.$$

Combining the above two equalities, we have

$$\begin{aligned} & \int_0^{1/e} u^{\kappa^{\gamma_n} + z - 1} (1 - u)^{\kappa - \kappa^{\gamma_n}} \ln \frac{1}{u} \, du \\ &= R_{\gamma_n}^{\kappa} \kappa^{-(1-\gamma_n)} \left\{ (1 - \gamma_n) \ln \kappa \left(1 + \mathcal{O}(\kappa^{-\gamma_n}) \right) \kappa^{-(1-\gamma_n)z} + \mathcal{O}(\kappa^{-(1-\gamma_n)z}) \right\}, \quad \kappa \xrightarrow{d} \infty. \end{aligned}$$

By substituting the last expression in (48) we reduce integral J_1 to the form

$$J_1 = \frac{R_{\gamma_n}^{\kappa}}{\Gamma(\varrho)} \kappa^{-(1-\gamma_n)} \left\{ (1 - \gamma_n) \ln \kappa \left(1 + \mathcal{O}(\kappa^{-\gamma_n}) \right) \int_0^{\zeta} \kappa^{-(1-\gamma_n)z} z^{\varrho-1} \, dz + \int_0^{\zeta} z^{\varrho-1} \mathcal{O}(\kappa^{-(1-\gamma_n)z}) \, dz \right\}. \quad (54)$$

Similarly to the above, we represent the integrals in (54) in terms of the incomplete Gamma function as

$$\int_0^{\zeta} \kappa^{-(1-\gamma_n)z} z^{\varrho-1} \, dz = \frac{\Gamma(\varrho)}{[(1 - \gamma_n) \ln \kappa]^{\varrho}} - \frac{\Gamma(\varrho, \zeta(1 - \gamma_n) \ln \kappa)}{[(1 - \gamma_n) \ln \kappa]^{\varrho}},$$

and use asymptotic expansion (52) to obtain

$$\int_0^{\zeta} \kappa^{-(1-\gamma_n)z} z^{\varrho-1} \, dz = \frac{\Gamma(\varrho)}{[(1 - \gamma_n) \ln \kappa]^{\varrho}} + \mathcal{O}\left(\frac{1}{\kappa^{\zeta(1-\gamma_n)} \ln \kappa}\right), \quad \kappa \xrightarrow{d} \infty.$$

Substituting this result in (54) and choosing ζ large enough, we find the asymptotic value of J_1 for $d \gg 1$:

$$J_1 = \frac{\Gamma(\kappa^{\gamma_n}) \Gamma(\kappa - \kappa^{\gamma_n} + 1)}{\Gamma(\kappa + 1)} (1 - \gamma_n)^{\beta} (\ln \kappa)^{\beta} \left(1 + \mathcal{O}\left(\frac{1}{\ln \kappa}\right) \right), \quad \kappa \xrightarrow{d} \infty.$$

Using (35), J_2 can be evaluated as

$$J_2 = \mathcal{O}\left(\frac{\Gamma(\kappa^{\gamma_n} + \zeta) \Gamma(\kappa - \kappa^{\gamma_n} + 1)}{\Gamma(\kappa + \zeta + 1)} \ln \kappa\right),$$

which finally yields the sought asymptotic representation of integral (39) for $d \rightarrow \infty$ and $\beta \in (0, 1)$:

$$\begin{aligned} & \int_0^1 u^{\alpha-1} (1 - u)^{\kappa-\alpha} \left(\ln \frac{1}{u}\right)^{\beta} \, du \\ &= \frac{\Gamma(\kappa^{\gamma_n}) \Gamma(\kappa - \kappa^{\gamma_n} + 1)}{\Gamma(\kappa + 1)} (1 - \gamma_n)^{\beta} (\ln \kappa)^{\beta} \left(1 + \mathcal{O}\left((\ln \kappa)^{-\beta}\right) \right), \quad \kappa \xrightarrow{d} \infty. \end{aligned} \quad (55)$$

Step 3. Now we use the developed expansions (53) and (55) to prove the unboundedness of the expected optimal value z^* of MAP when $F^{-1}(\cdot)$ has a logarithmic singularity (37) with $0 < \beta < 1$. Upon substitution of the asymptotic relation (37) into integral $I_{\alpha, \kappa}$ (13) one has

$$I_{\alpha, \kappa} \sim -v \int_0^1 u^{\alpha-1} (1 - u)^{\kappa-\alpha} \left(\ln \frac{1}{u}\right)^{\beta} \, du, \quad \kappa \xrightarrow{n, d} \infty.$$

Using the derived expressions (53) and (55) for the above integral, the upper bound \bar{z}^* of the expected optimal value z^* of the MAP can be asymptotically evaluated as

$$\bar{Z}_{d,n}^* = -v\varpi_n n(\ln \kappa)^\beta + \mathcal{O}(n), \quad \kappa \xrightarrow{n,d} \infty,$$

where $\varpi_n = 1$ for $\kappa \xrightarrow{n} \infty$, and $\varpi_n = (1 - \gamma_n)^\beta$ if $\kappa \xrightarrow{d} \infty$. From the last equality it immediately follows that $\bar{Z}_{d,n}^*$ is unbounded from below for large n and d , which validates the assertion (38) of the theorem: $\lim_{d \rightarrow \infty} Z_{d,n}^* = \lim_{n \rightarrow \infty} Z_{d,n}^* = -\infty$ for $0 < \beta < 1$.

Step 4. To extend the last result to values of $\beta \geq 1$ in (37), we recall that by virtue of decomposition (40) the main contribution to the asymptotic value of integral (39) is made by the first integral in the right-hand side of (40). For any $\beta \geq 1$, this integral is majorized as

$$-v \int_0^{1/e} u^{\alpha-1} (1-u)^{\kappa-\alpha} \left(\ln \frac{1}{u}\right)^\beta du \leq -v \int_0^{1/e} u^{\alpha-1} (1-u)^{\kappa-\alpha} \left(\ln \frac{1}{u}\right)^{\beta_0} du, \quad (56)$$

where $0 < \beta_0 < 1$. From the above results it follows that the right-hand side of (56) approaches $-\infty$ when $\kappa \xrightarrow{n,d} \infty$. Thus, we conclude that $\bar{Z}_{d,n}^* \rightarrow -\infty$, and, consequently, $Z_{d,n}^* \rightarrow -\infty$ when $\kappa \xrightarrow{n,d} \infty$ and $\beta \geq 1$ in (37). \blacksquare

It is easy to see that Theorems 2 and 3 can be applied to deal with situations when $F^{-1}(x)$ has a singularity of the form $F^{-1}(x) \sim -vx^{-\beta_1} \left(\ln \frac{1}{x}\right)^{\beta_2}$, $x \rightarrow 0+$, where $v, \beta_2 > 0$, and $0 < \beta_1 < 1$. In such a case, again, one has $\lim_{d \rightarrow \infty} Z_{d,n}^* = \lim_{n \rightarrow \infty} Z_{d,n}^* = -\infty$.

To finalize the presented analysis of the expected optimal value of the MAP, we consider the special case of a random MAP (1) with $n = 2$, which has been omitted from the preceding discussion. Recall that according to Lemma 3, $\gamma_n < 1/2$ for all $n \geq 3$, whereas a simple calculation confirms that $\gamma_2 = 1/2$. This precludes a direct application of the arguments of Lemma 4 and Theorems 1–3 to the case of $n = 2$, since terms of type $\mathcal{O}(\kappa^{2\gamma_n-1})$ do not vanish for $n = 2$ (see, for example, the proof of Lemma 4, etc).

This ramification is easily circumvented by means of representation (11), which allows us to rewrite the expected optimal cost $Z_{d,2}^*$ of a $n = 2$ random MAP in a form similar to expression (4) for the lower bound $\underline{Z}_{d,n}^*$, namely $Z_{d,2}^* = \mathbb{E}_{F_2}[X_{(1/\kappa)}]$. Here $\kappa = 2^{d-1}$ and $F_2 = F * F$ is the distribution of sum of two independent F -distributed random variables. The interior of the support of F_2 has the form $(2a, 2b)$, where the usual convention

$$n \cdot (\pm\infty) = \pm\infty, \quad n \in \mathbb{N} \setminus \{0\} \quad (57)$$

is adopted. Then, the reasonings of Lemma 4 and Theorem 1 can be readily employed to show that the expected optimal value of a random $n = 2$ MAP satisfies $\lim_{d \rightarrow \infty} Z_{d,2}^* = 2a$ if $a \in \mathbb{R}$ and $F_2^{-1}(\cdot)$ has an asymptotic expansion of form (24): $F_2^{-1}(u) = 2a + \mathcal{O}(u^\beta)$ for some $\beta > 0$ as $u \rightarrow 0+$. Similarly, it can be demonstrated that $\lim_{d \rightarrow \infty} Z_{d,2}^* = -\infty$ when $a = -\infty$ and $F_2^{-1}(\cdot)$ conforms to conditions (27) or (37).

To merge these observations with the statements of Theorems 1–3 (where conditions are imposed on F^{-1} , not F_2^{-1}), it only remains to verify that $F_2^{-1}(\cdot)$ satisfies conditions (24), (27), or (37) as soon as $F^{-1}(\cdot)$ does. Indeed, in the case of a finite a , the convolution representation of F_2 produces for $x \ll 1$

$$F_2(2a+x) = \int_a^{a+x} F(a+x-(y-a)) dF(y) \leq F(a+x) \int_a^{a+x} dF(y) = F^2(a+x),$$

and

$$F_2(2a + x) \geq \int_a^{a+\frac{x}{2}} F(a + x - (y - a)) dF(y) \geq F(a + \frac{x}{2}) \int_a^{a+\frac{x}{2}} dF(y) = F^2(a + \frac{x}{2}),$$

which by inversion implies that $F^{-1}(\sqrt{x}) - a \leq F_2^{-1}(x) - 2a \leq 2(F^{-1}(\sqrt{x}) - a)$ for a sufficiently small positive x . Dividing the last inequalities by $x^{\beta/2}$ and taking into account (24) we obtain that the ratio $(F_2^{-1}(x) - 2a)/x^{\beta/2}$ is bounded for $x \rightarrow 0+$, which means that $F_2^{-1}(x)$ is equal to $2a + \mathcal{O}(x^{\beta/2})$ in the vicinity of $x = 0+$, and therefore complies with the conditions of Theorem 1.

In the case of $a = -\infty$, the function $F_2(x)$ for $-x \gg 1$ can be bounded from above as

$$F_2(2x) = \left(\int_{-\infty}^x + \int_x^b \right) F(x - (y - x)) dF(y) \leq \int_{-\infty}^x 1 \cdot dF(y) + F(x) \int_x^b dF(y) \leq 2F(x),$$

whereas a lower bound is computable in the form

$$F_2(x) \geq \int_{-\infty}^x F(x - y) dF(y) \geq F(0)F(x).$$

Inverting the derived bounds, we have $2F^{-1}(\frac{x}{2}) \leq F_2^{-1}(x) \leq F^{-1}(\frac{x}{F(0)})$, $x \rightarrow 0+$. Applying the same reasoning as above, we obtain that $F_2^{-1}(x)$ has an asymptotical expansion of the form (27) for $x \rightarrow 0+$, provided that $F^{-1}(\cdot)$ satisfies (27). If $F^{-1}(\cdot)$ has asymptotic representation (37) in the vicinity of 0, the derived bounds on $F_2^{-1}(\cdot)$ can be used to show that $\lim_{d \rightarrow \infty} Z_{d,2}^* = -\infty$ in this case as well.

Using the notation (57), we can summarize the developed results in the following theorem.

Theorem 4 (Expected Optimal Value of the MAP) *Consider a $d \geq 3, n \geq 2$ MAP (1) with cost coefficients that are iid random variables from an absolutely continuous distribution F with existing first moment. Let (a, b) , where a and b are finite or infinite, be the interior of the support set of this distribution. If the distribution F satisfies conditions of either of Theorems 1, 2, or 3, the expected optimal value of the MAP satisfies*

$$Z_{d,n}^* \rightarrow na, \quad n \rightarrow \infty \text{ or } d \rightarrow \infty.$$

The foregoing analysis is constructive in the sense that it allows for deriving practically useful bounds for the expected optimal value of random MAP. The next section discusses this issue in detail and presents experimental evidence on the goodness of the developed bounds.

4 Asymptotical bounds and rate of convergence for the expected optimal value of MAP

Although the primary focus of the preceding sections has been on determining the limiting value of the expected optimal cost of “infinitely large” random MAPs, the obtained results can be readily employed to construct upper and lower asymptotical bounds for the expected optimal value of MAP when one of the parameters n or d is large but finite. The lower and upper bounds $Z_{d,n}^*$ and $\bar{Z}_{d,n}^*$ introduced in Lemma 2 are especially useful when the support of distribution F of the MAP’s cost coefficients has a finite left endpoint. The following statement is a special case of Lemma 4 and Theorem 1.

Corollary 1 Consider a $d \geq 3$, $n \geq 3$ MAP (1) with cost coefficients that are iid random variables from an absolutely continuous distribution with existing first moment. Let (a, b) , where $a \in \mathbb{R}$, $b \in (a, +\infty]$, be the interior of the support set of this distribution, and assume that the inverse $F^{-1}(u)$ of the c.d.f. $F(u)$ of the distribution is such that

$$F^{-1}(u) \sim a + \sum_{s=1}^{\infty} a_s u^{s/\mu}, \quad u \rightarrow 0+, \quad \mu > 0. \quad (58)$$

Then, for any integer $m \geq 1$, lower and upper bounds $\underline{Z}_{d,n}^*$, $\overline{Z}_{d,n}^*$ (4) on the expected optimal cost $Z_{d,n}^*$ of the MAP can be asymptotically evaluated as

$$\underline{Z}_{d,n}^* = an + \sum_{s=1}^{m-1} a_s \frac{n \Gamma(\kappa + 1) \Gamma(\frac{s}{\mu} + 1)}{\Gamma(\kappa + \frac{s}{\mu} + 1)} + \mathcal{O}\left(n \frac{\Gamma(\kappa + 1) \Gamma(\frac{m}{\mu} + 1)}{\Gamma(\kappa + \frac{m}{\mu} + 1)}\right), \quad \kappa \xrightarrow{n,d} \infty, \quad (59a)$$

$$\overline{Z}_{d,n}^* = an + \sum_{s=1}^{m-1} a_s \frac{n \Gamma(\kappa + 1) \Gamma(\frac{s}{\mu} + \alpha)}{\Gamma(\alpha) \Gamma(\kappa + \frac{s}{\mu} + 1)} + \mathcal{O}\left(n \frac{\Gamma(\kappa + 1) \Gamma(\frac{m}{\mu} + \alpha)}{\Gamma(\alpha) \Gamma(\kappa + \frac{m}{\mu} + 1)}\right), \quad \kappa \xrightarrow{n,d} \infty. \quad (59b)$$

It can be shown that the lower and upper bounds defined by (59) are convergent, i.e.,

$$|\overline{Z}_{d,n}^* - \underline{Z}_{d,n}^*| \rightarrow 0, \quad \kappa \xrightarrow{n,d} \infty,$$

whereas the corresponding asymptotical bounds for the case of distributions with support unbounded from below may be divergent in the sense that $|\overline{Z}_{d,n}^* - \underline{Z}_{d,n}^*| \not\rightarrow 0$ when $\kappa \xrightarrow{n,d} \infty$.

Asymptotical representations (59) for the bounds $\underline{Z}_{d,n}^*$ and $\overline{Z}_{d,n}^*$ are simplified when the inverse $F^{-1}(\cdot)$ of the c.d.f. of the distribution has a regular power series expansion in the vicinity of zero. Assume, for example, that function $F^{-1}(\cdot)$ can be written as

$$F^{-1}(u) = a + a_1 u + \mathcal{O}(u^2), \quad u \rightarrow 0+. \quad (60)$$

The asymptotic expressions (59) for the lower and upper bounds $\underline{Z}_{d,n}^*$ and $\overline{Z}_{d,n}^*$ for the expected optimal cost $Z_{d,n}^*$ then take the form

$$\underline{Z}_{d,n}^* \sim na + \frac{a_1}{n^{d-2}}, \quad \kappa \xrightarrow{n,d} \infty, \quad (61a)$$

$$\overline{Z}_{d,n}^* \sim na + a_1 \frac{[e^{d-1}]}{n^{d-2}}, \quad \kappa \xrightarrow{n} \infty, \quad (61b)$$

$$\overline{Z}_{d,n}^* \sim na + a_1 n^{\gamma_n^{(d-1)-(d-2)}}, \quad \kappa \xrightarrow{d} \infty. \quad (61c)$$

It is of interest to note that if $a = 0$ in (60) then for $n \gg 1$ and d fixed the expected optimal value of the MAP is asymptotically bounded as

$$\frac{a_1}{n^{d-2}} + \mathcal{O}\left(\frac{1}{n^{d-1}}\right) \leq Z_{d,n}^* \leq \frac{a_1 [e^{d-1}]}{n^{d-2}} + \mathcal{O}\left(\frac{1}{n^{d-1}}\right), \quad n \rightarrow \infty, \quad (62)$$

which immediately yields the rate of convergence to zero for $Z_{d,n}^*$ as n approaches infinity:

Corollary 2 Consider a $d \geq 3$, $n \geq 3$ MAP (1) with cost coefficients that are iid random variables from an absolutely continuous distribution with existing first moment. Let $(0, b)$, where $b \in (0, +\infty]$, be the interior of the support set of this distribution, and assume that the inverse $F^{-1}(u)$ of the c.d.f. $F(u)$ of the distribution satisfies (60) with $a = 0$. Then, for a fixed d and $n \rightarrow \infty$ the expected optimal value $Z_{d,n}^*$ of the MAP converges to zero as $\mathcal{O}(n^{-(d-2)})$.

Furthermore, inequalities (62) allow one to conjecture that the expected optimal value of a random MAP with an absolutely continuous distribution F , which satisfies (60) with $a = 0$, has the form

$$Z_{d,n}^* = \frac{C}{n^{d-2}} + \mathcal{O}\left(\frac{1}{n^{d-1}}\right), \quad n \gg 1,$$

where constant $C > 0$ is such that $a_1 \leq C \leq a_1 \lceil e^{d-1} \rceil$.

We illustrate the tightness of the developed bounds (59) by comparing them to the computed expected optimal values of MAPs with coefficients c_{i_1, \dots, i_d} drawn from the uniform $U(0, 1)$ distribution and exponential distribution with mean 1. The p.d.f. of the uniform distribution is given by $f_{U(0,1)}(x) = \mathbf{1}_{[0,1]}(x)$, and that of exponential with mean 1 distribution is $f_{\text{expo}(1)}(x) = e^{-x} \mathbf{1}_{[0,+\infty)}(x)$, where $\mathbf{1}_A(x)$ is the indicator function of a set $A \subseteq \mathbb{R}$. It is elementary that the inverse functions $F^{-1}(\cdot)$ of the c.d.f.'s for both these distributions are representable in the form (60) with $a = 0$, $a_1 = 1$, thereby reducing the general expressions (59) for the lower and upper bounds $\underline{Z}_{d,n}^*$ and $\overline{Z}_{d,n}^*$ to the form (61).




Figure 1: Expected optimal value $Z_{d,n}^*$, lower and upper bounds $\underline{Z}_{d,n}^*$, $\overline{Z}_{d,n}^*$ of an MAP with fixed $d = 3$ (left) and $d = 5$ (right) for uniform $U(0, 1)$ and $\text{exponential}(1)$ distributions.

The numerical experiments involved solving multiple instances of randomly generated MAPs with the number of dimensions d ranging from 3 to 10, and the number n of elements in each dimension running from 3 to 20. The number of instances generated for estimation of the expected optimal value of the MAP with a given distribution of cost coefficients varied from 1000 (for smaller values of d and n) to 50 (for problems with largest n and d).

To solve the problems to optimality, we used a branch-and-bound algorithm that navigated through the index tree representation of the MAP. Figures 1 and 2 display the obtained expected optimal values of MAP with uniform and exponential iid cost coefficients when d is fixed at $d = 3$ or 5 and $n = 3, \dots, 20$, and when $n = 3$ or 5 and d runs from 3 to 10. This ‘‘asymmetry’’ in reporting of the results is explained by the fact that the implemented branch-and-bound algorithm based on index tree is more efficient in




Figure 2: Expected optimal value $Z^*_{d,n}$, lower and upper bounds $\underline{Z}^*_{d,n}$, $\bar{Z}^*_{d,n}$ of an MAP with fixed $n = 3$ (left) and $n = 5$ (right) for uniform $U(0, 1)$ and $\text{exponential}(1)$ distributions.

solving “shallow” MAPs, i.e., instances that have larger n and smaller d . The solution times varied from several seconds to 20 hours on a 2GHz PC.

Figures 1 and 2 suggest that MAPs with exponentially and uniformly distributed costs and the same dimensionality possess very close expected optimal values. This observation is in complete agreement with the presented above theoretical findings: indeed, we argue that the asymptotical behavior of large-scale random MAPs is determined by the properties of the inverse $F^{-1}(x)$ of the c.d.f. $F(\cdot)$ at $x = 0$. In the considered case of uniformly and exponentially distributed random MAPs, the inverse functions $F^{-1}(\cdot)$ of c.d.f.’s of these two distributions share the first two terms of their asymptotical expansions at zero, which leads to similar numerical values of the expected optimal cost.

The conducted numerical experiments also suggest that the constructed lower and upper bounds for the expected optimal cost of random MAPs are quite tight, with the upper bound $\bar{Z}^*_{d,n}$ being tighter for the case of fixed n and large d (see Figs. 1–2).

Conclusions

In this paper we have conducted asymptotical analysis of the expected optimal value of the Multidimensional Assignment Problem where the assignment costs are iid random variables drawn from a continuous distribution. It has been demonstrated that asymptotical behavior of the expected optimal cost of a random MAP in the case when one of the problem’s dimension parameters approaches infinity is determined by the location of the left endpoint a of the support $[a, b]$ of the distribution. Namely, the expected optimal cost of a d -dimensional random MAP with n elements in each dimension approaches the limiting value of na when one of the parameters n or d is fixed and the other increases infinitely. This statement has been proven to hold true for a broad class of continuous distributions with a finite or infinite left endpoint of the support. The presented analysis is constructive in the sense that it allows for derivation of lower and upper asymptotical bounds for the expected optimal value of the problem for a prescribed probability distribution. If the distribution’s support set has a finite left endpoint, these bounds are converging.

References

- [1] Aiex, R. M., Resende, M. G. C., Pardalos, P. M., and G. Toraldo (2005) GRASP with Path Relinking for Three-Index Assignment, *INFORMS Journal on Computing*, 17, in press.
- [2] Aldous, D., (1992) Asymptotics in the random assignment problem, *Probability Theory and Related Fields*, 93, 507–534.
- [3] Aldous, D., (2001) The $\zeta(2)$ limit in the random assignment problem, *Random Structures and Algorithms*, 18(4) 381–418.
- [4] Alon, N., and J. Spencer (2000) *The Probabilistic Method*, 2nd edition, Interscience Series in Discrete Mathematics and Optimization, John Wiley, New York.
- [5] Andrijich, S. M., and L. Caccetta (2001) Solving the multisensor data association problem, *Nonlinear Analysis*, 47, 5525–5536.
- [6] Balas, E., and M. J. Saltzman (1991) An algorithm for the three-index assignment problem, *Operations Research*, 39, 150–161.
- [7] Burkard, R. E. (2002) Selected topics on assignment problems, *Discrete Applied Mathematics*, 123, 257–302.
- [8] Burkard, R. E., and E. Çela (1999) Linear assignment problems and extensions, *In: D.-Z. Du and P. M. Pardalos (Eds.), Handbook of Combinatorial Optimization*, Vol. 4, Kluwer Academic Publishers, Dordrecht.
- [9] Burkard, R. E., and E. Çela (1997) Quadratic and three-dimensional assignments, *In: M. DellAmico, F. MaQoli, and S. Martello (Eds.) Annotated Bibliographies in Combinatorial Optimization*, Wiley, Chichester, 373–391.
- [10] Clemons, W., Grundel, D., and D. Jeffcoat (2003) Applying simulated annealing on the multidimensional assignment problem, *International Conference on Cooperative Control and Optimization*, proceedings.
- [11] David, H. A. (1981) *Order Statistics*, 2nd ed. New York, Wiley.
- [12] Erdélyi, A. (1961) General Asymptotic Expansions of Laplace Integrals, *Archives of Rational Mechanics Analysis*, 7, 1–20.
- [13] Erdélyi, A. (Ed.) (1953) *Higher Transcendental Functions*, vol. 1, McGraw-Hill Book Company, New York.
- [14] Garey, M. R., and D. S. Johnson (1979) *Computers and Intractability: A Guide to the Theory of NP-completeness*, W. H. Freeman and Company, 1979.
- [15] Goemans M. X., and M. Kodilian (1993) A lower bound on the expected value of an optimal assignment, *Mathematics of Operations Research*, 18, 267–274.
- [16] Grundel, D. A., Oliveira, C. A. S., and P. M. Pardalos (2004) Asymptotic properties of random multidimensional assignment problems, *Journal of Optimization Theory and Applications*, 122(3), 487–500.

- [17] Grundel, D. A., Krokhmal, P. A., Oliveira, C. A. S., and P. M. Pardalos (2004) On the Number of Local Minima in the Multidimensional Assignment Problem, *Research Report #2004-8*, ISE Dept., University of Florida, available at <http://www.engineering.uiowa.edu/~krokhmal/pdf/map-localmins.pdf>
- [18] Karp, R. M. (1987) An upper bound on the expected cost of an optimal assignment, *Discrete Algorithms and Complexity*, Academic Press, Boston, 1–4.
- [19] Lazarus, A. J. (1990) Certain expected values in the random assignment problem, *Technical Report*, Math. & Computer Science Dept., Univ of California, Riverside.
- [20] Lazarus, A. J. (1993) Certain expected values in the random assignment problem, *Operations Research Letters*, 14, 207–214.
- [21] Linusson, S., and J. Wästlund (2003) A proof of Parisi’s conjecture on the random assignment problem, *Working paper*.
- [22] Mézard, M., and G. Parisi (1985) Replicas and optimization, *Journal de Physique Lettres*, 46(17), 771–778.
- [23] Mézard, M., and G. Parisi (1987) On the solution of the random link matching problems, *Journal de Physique*, 48(9), 1451–1459.
- [24] Murphey, R., Pardalos, P., and L. Pitsoulis (1998a) A greedy randomized adaptive search procedure for the multitarget multisensor tracking problem, *In: DIMACS Series Vol. 40*, American Mathematical Society, 277–302.
- [25] Murphey, R., Pardalos, P. M., and L. S. Pitsoulis (1998b) A Parallel GRASP for the Data Association Multidimensional Assignment Problem, *In: Parallel Processing of Discrete Problems*, The IMA Volumes in Mathematics and its Applications, Vol. 106, Springer, Berlin, 159–180.
- [26] Nair, C., Prabhakar, B., and M. Sharma (2003) A Proof of the Conjecture due to Parisi for the Finite Random Assignment Problem, *Working Paper*.
- [27] Olin, B. (1992) *Asymptotic properties of the random assignment problem*, Ph.D. thesis, Division of Optimization and Systems Theory, Department of Mathematics, Royal Institute of Technology, Stockholm, Sweden.
- [28] Olver, F. W. (1997) *Asymptotics and Special Functions*, 2nd edition, AK Peters Ltd, Wellesley, MA.
- [29] Pardalos, P. and Pitsoulis, L., Eds. (2000) *Nonlinear Assignment: Problems, Algorithms and Applications*, Kluwer Academic Publishers, Dordrecht.
- [30] Pardalos, P. M., and K. G. Ramakrishnan (1993) On the expected optimal value of random assignment problems: Experimental results and open questions, *Computational Optimization and Applications*, 2, 261–271.
- [31] Parisi, G. (1998) A conjecture on random bipartite matching, *Physics e-Print archive*, <http://xxx.lang.gov/ps/cond-mat/9801176>.

- [32] Papadimitrou, C. H., and K. Steiglitz (1998) *Combinatorial Optimization: Algorithms and Complexity*, Dover, New York.
- [33] Pasiliao, E. L. (2003) *Algorithms for Multidimensional Assignment Problems*, PhD thesis, Department of Industrial and Systems Engineering, University of Florida.
- [34] Pierskalla, W. (1968). The multidimensional assignment problem, *Operations Research*, 16, 422–431.
- [35] Poore, A. B. (1994a) Multidimensional assignment formulation of data association problems arising from multitarget and multisensor tracking, *Computation Optimization and Applications*, 3, 27–54.
- [36] Poore, A. B. (1994b) A numerical study of some data association problems arising in multitarget tracking, *In: W. W. Hager, D. W. Hearn, P. M. Pardalos (Eds.) Large Scale Optimization: State of the Art*, Kluwer Academic Publishers, Dordrecht, 339–361.
- [37] Poore, A. B., Rijavec, N., Liggins, M., and V. Vannicola (1993) Data association problems posed as multidimensional assignment problems: problem formulation, *In: O. E. Drummond (Ed.) Signal and Data Processing of Small Targets*, SPIE, Bellingham, WA, 552–561.
- [38] Pusztaszeri, J., Rensing, P. E., and T. M. Liebling (1995) Tracking elementary particles near their primary vertex: a combinatorial approach, *Journal of Global Optimization*, 16, 422–431.
- [39] Veenman, C. J., Hendriks, E. A., and M. J. T. Reinders (1998) A fast and robust point tracking algorithm, *Proceedings of the Fifth IEEE International Conference on Image Processing*, 653–657, Chicago, USA.
- [40] Walkup, D. W. (1979) On the expected value of a random assignment problem, *SIAM Journal on Computation*, 8, 440–442.
- [41] Wolfram Research, Inc. <http://functions.wolfram.com>